

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 659 884 A2**

(12)

**EUROPEAN PATENT APPLICATION**(21) Application number: **94120679.9**(51) Int. Cl.<sup>6</sup>: **C12N 15/29, C12N 1/21,  
C07K 14/415**(22) Date of filing: **27.12.94**

(83) Declaration under Rule 28(4) EPC (expert solution)

(30) Priority: **24.12.93 JP 327943/93**(43) Date of publication of application:  
**28.06.95 Bulletin 95/26**(84) Designated Contracting States:  
**DE FR GB**

(71) Applicant: **SUMITOMO CHEMICAL COMPANY,  
LIMITED**  
**5-33 Kitahama 4-chome**  
**Chuo-ku**  
**Osaka-shi**  
**Osaka (JP)**

(72) Inventor: **Yamamoto, Mika**  
**Gurandosezon 306,**  
**2-21, Yamata-cho**  
**Sakado-shi,**  
**Saitama-ken (JP)**  
Inventor: **Oheda, Kenji, Pereniarufulakksu 1,6**  
**Matsugasakikudo-cho,**  
**Sakyo-ku**  
**Kyoto-shi,**  
**Kyoto-fu (JP)**

(74) Representative: **VOSSIUS & PARTNER**  
**Postfach 86 07 67**  
**D-81634 München (DE)**

(54) **Carrot 16 kd protein, gene coding for said protein and plasmid containing said gene.**

(57) There are disclosed a protein corresponding to a molecular weight of approximately 16 kD, which can be specifically expressed in carrot roots, a gene coding for the protein, and a plasmid containing the gene.

**EP 0 659 884 A2**

The present invention relates to a carrot 16 kD protein, and more particularly, it relates to a protein corresponding to a molecular weight of approximately 16 kD, which can be specifically expressed in carrot roots. The present invention further relates to a gene coding for said protein and a plasmid containing said gene.

It is well known that reserve proteins are present in the subterranean storage tissue of sweet potatoes, white potatoes and the like. By contrast, the presence of reserve proteins has not yet been found in the subterranean tissue of root vegetables which are crops having edible roots such as carrots.

Carrot roots have been used as a root vegetable for food; it has, however, been found that they contain only small amounts of proteins essential to the maintenance of health. Thus, it has been desired that high-protein carrots are developed by higher expression of proteins which are present in small amounts but specifically in the carrot roots. For this purpose, it has been necessary to find, first of all, proteins which can be specifically expressed in carrot roots and genes coding for these proteins.

Under these circumstances, the present inventors have intensively studied. As a result, they have found a soluble protein which can be specifically expressed in carrot roots, and succeeded in the cloning of cDNA coding for said protein. Further, they have determined the amino acid sequence of said protein by analyzing the base sequence of said cDNA, thereby completing the present invention.

Thus, the present invention provides: a protein corresponding to a molecular weight of approximately 16 kD, which can be specifically expressed in carrot roots, and having the amino acid sequence as shown in Sequence Listing, SEQ ID NO: 1; a gene containing a coding region for the amino acid sequence of said protein; and a plasmid containing said gene.

Figure 1 shows the results of the SDS-polyacrylamide electrophoresis (with silver staining) which was performed with all the proteins present in carrot roots as a sample. Lane M is corresponding to a molecular weight marker; and lane 1, the sample. The arrow indicates the band corresponding to the protein of the present invention.

Figure 2 shows the N-terminal amino acid sequence (one-character representation) of the protein of the present invention. This sequence is corresponding to the amino acids at the 2nd to the 36th positions in the amino acid sequence as shown in Sequence Listing, SEQ ID NO: 1.

Figure 3 shows the structure of plasmid pC16 containing the gene coding for the protein of the present invention.

Figure 4 shows the results of the SDS-polyacrylamide electrophoresis (with silver staining) which was performed to examine the tissue specificity of the protein of the present invention. Lane M is corresponding to a molecular weight marker; lane 1, the sample of all the proteins present in the entire plant on the 4th week after the seeding; lane 2, the sample of all the proteins present in the leaves on the 7th week after the seeding; lane 3, the sample of all the proteins present in the roots on the 7th week after the seeding; lane 4, the sample of all the proteins present in the leaves on the 9th week after the seeding; lane 5, the sample of all the proteins present in the roots on the 9th week after the seeding; and lane 6, the sample of all the proteins on the 21st week after the seeding. The arrow indicates the band corresponding to the protein of the present invention.

Figure 5 shows the results of the SDS-polyacrylamide electrophoresis (with silver staining) which was performed to examine the comparison between the varieties of the protein of the present invention. Lane M is corresponding to a molecular weight marker; lane 1, the sample of all the protein present in the root of carrot "Kuroda Gosun"; lane 2, the sample of all the proteins present in the root of carrot "Early-chantenee"; lane 3, the sample of all the proteins present in the root of carrot "Imperator"; and lane 4, the sample of all the proteins present in the root of carrot "Nuntesscarlet". The arrow indicates the band corresponding to the protein of the present invention.

Figure 6 shows the results of the search on the data bases EMBL and NBRF to examine the homology of the protein of the present invention.

As used herein, the term "gene" refers to all the corresponding nucleic acids with genetic information, including genomic DNAs, their corresponding mRNAs and their corresponding cDNAs. The gene of the present invention is to be construed to include genes each containing a coding region for the amino acid sequence as shown in Sequence Listing, SEQ ID NO: 1, but each corresponding to another base sequence than the base sequence as shown Sequence Listing, SEQ ID NO: 2.

The protein of the present invention can be widely found in many varieties of carrots (*Daucus carota* L.), such as "Kuroda Gosun", "Early Chantenee", "Imperator" and "Nuntesscarlet", and it is specifically present in their roots. The protein of the present invention has a molecular weight of approximately 16 kD (the calculated value based on the amino acid sequence is 16,125.03). The protein of the present invention has the amino acid sequence as shown in Sequence Listing, SEQ ID NO: 1, which is further corresponding to, for example, the base sequence as shown in Sequence Listing, SEQ ID NO: 2. The utilization of a gene

coding for the amino acid sequence of such a protein of the present invention makes it possible to develop high-protein carrots, and the introduction and expression of said gene into other root vegetable crops such as radishes, turnips, sugar beets and burdocks make it possible to perform the breeding of these crops for increasing the amount of proteins. Further, the search on data bases such as EMBL and NBRF for genes and amino acid sequences, which have a high degree of homology with the base sequence coding for the protein of the present invention and the corresponding amino acid sequence, respectively, makes it possible to attain, for examples, the following matters: (1) when they have a high degree of homology with the gene and amino acid sequence of a certain protein related to disease resistance, the breeding for imparting a resistance to disease and the elucidation of a mechanism of the disease resistance can be attained by ordinary techniques for genetic engineering using the gene containing a coding region for the amino acid sequence of the protein of the present invention; (2) when they have a high degree of homology with the gene and amino acid sequence of a certain protein expressed and derived in response to the change of various plant hormones, various kinds of breeding which make use of stress response or hormone response can be attained by ordinary techniques for genetic engineering using the gene containing a coding region for the amino acid sequence of the protein of the present invention; (3) when they have a high degree of homology with the gene and amino acid sequence of a certain protein responsible for pollen allergy, the breeding of non-allergic varieties with an improvement of pollen and a reduction in the amount of pollen expressed can be attained by ordinary techniques for genetic engineering using the gene containing a coding region for the amino acid sequence of the protein of the present invention; and (4) when they have a high degree of homology with the gene and amino acid sequence of a certain heat-shock protein, the improvement in the storage of useful proteins thorough the support of their holding and the effective utilization for the breeding to improve the transport of useful proteins can be attained by ordinary techniques for genetic engineering using the gene containing a coding region for the amino acid sequence of the protein of the present invention.

The protein of the present invention is present in the root tissues of many varieties of commercially available carrots, such as "Kuroda Gosun", "Early Chantenee", "Imperator" and "Nuntesscaret", and it is one of the soluble proteins which can be extracted from these root tissues.

For the separation, purification or detection of the protein of the present invention, for example, the following method can be used.

First of all, root tissues are thoroughly ground with an ordinary buffer used for soluble-protein extraction, such as a solution of potassium phosphate or sodium borate, by a commercially available mixer or a whirling blender, and the tissue residue is removed by filtration with gauze or by centrifugation at 10,000 rpm for about 10 minutes to give a crude extract of the protein of the present invention. The above buffer for soluble-protein extraction may optionally contain a reducing agent such as ascorbic acid or 2-mercaptoethanol.

The amount of the buffer for soluble-protein extraction to be used for the extraction of the protein of the present invention from the root tissues can be, for example, about 10 ml to about 100 ml, preferably about 10 ml to about 25 ml, per 5 g of the root tissues.

The above crude extract of the protein of the present invention can be suitably concentrated by ultrafiltration with Centriprep-10 (Amicon) or the like.

For the separation and purification of the protein of the present invention from the above crude extract on the basis of a difference in molecular weight, there can be used an ordinary method such as SDS-polyacrylamide gel electrophoresis (U.K. Laemmli, *Nature*, 227,680(1970)) or liquid chromatography through gel filtration. The use of such a method results in that the protein of the present invention is separated to have a size of approximately 16 kD in molecular weight. Further, the protein of the present invention can be simply detected by an immunological method using a polyclonal or monoclonal antibody which is prepared by using, as an antigen, a fraction separated and purified by liquid chromatography through gel filtration or a fraction cut out after the separation in a gel by SDS-polyacrylamide gel electrophoresis.

The gene of the present invention can be obtained according to an ordinary method for genetic engineering as described in, for example, J. Sambrook, E.F. Frisch, & T. Maniatis, *Molecular Cloning*, 2nd ed., Cold Spring Harbor Laboratory, 1989.

This method will hereinafter be explained in detail. First of all, the entire RNA is extracted by an ordinary method such as guanidine hydrochloride/phenol method, SDS-phenol method or guanidine thiocyanate/CsCl method, in which the plant tissues of various plants are ground in a solution containing a strong protein-denaturing agent such as guanidine hydrochloride or guanidine thiocyanate, and proteins and other contaminants are removed by centrifugation or by phenol or chloroform extraction, after which the entire RNA is collected by ethanol precipitation. As a kit based on these methods, for example, ISOGEN (Nippon Gene) and Extraction-A-PLANT™ RNA ISOLATION KIT (Clontech) are commercially available. For

the purification of only mRNA from the entire RNA obtained, affinity chromatography utilizing hybridization between the poly-A tail of the mRNA and the oligo-dT tract combined with cellulose is effective. Oligo-dT cellulose is commercially available from Pharmacia or Collaborative. The poly-A tail-containing mRNA extracted from the column is collected by ethanol precipitation. As a method for the synthesis of cDNA from this poly-A tail-containing mRNA, there can be mentioned, for example, ordinary methods in which (1) the mRNA is used as a template and an oligo-dT primer is annealed to the poly-A tail thereof, after which first-strand DNA is synthesized by reverse transcriptase, followed by insertion of nicks and gaps in the RNA sequence using *Escherichia coli* RNaseH; the fragments of the RNA are then used as the subsequent primers to synthesize double-strand DNA using *E. coli* DNA polymerase I; (2) further, both ends are made blunt by T4 DNA polymerase, after which the resulting cDNA is purified and collected by phenol or chloroform extraction or by ethanol precipitation (as described in U. Gubler & B.J. Hoffman, *Gene*, 25, 263- (1983); H. Okayama & P. Berg, *Mol. Cell. Biol.*, 2, 161(1982)). As a kit based on these methods, for example, cDNA synthesis system PLUS (Amersham) is commercially available.

The cDNA thus obtained is ligated to an appropriate linker and an appropriate adaptor at both blunt ends using T4 DNA ligase, and excess free linkers and adapters are separated by gel filtration or the like, followed by ligation to an appropriate vector arm such as a  $\lambda$  phage-derived vector. Further, phage particles formed by *in vitro* packaging are inoculated into *E. coli* cells to form a plaque on the medium. In this manner, a cDNA library can be prepared.

For the selection of cDNA clones containing a coding region for the amino acid sequence of the protein of the present invention from the cDNA library, for example, immunoscreening with an antibody against the protein of the present invention is effective in a vector system such as  $\lambda$ gt 11, or plaque hybridization in which an oligonucleotide having the base sequence presumed from the amino acid sequence of the protein of the present invention is synthesized and labelled with a radioactive isotope (RI) or fluorescent reagent to use as a probe is effective in a vector system such as  $\lambda$ gt 10. Further, screening can also be effected by purifying DNA of the cDNA-containing phage and then conducting southern blot hybridization using the DNA as a probe.

The above probe can be produced by an ordinary method in which the number and sequence of amino acids at the N-terminal side are determined by the analysis of the purified protein of the present invention with an amino acid sequencer based on the Edman degradation (e.g., Amino Acid Sequencer 473A, Applied Biosynthesis), and the information on the amino acid sequence is used for substitution with a series of base codons presumed therefrom.

The cDNA clone thus obtained, which contains a coding region for the amino acid sequence of the protein of the present invention, is subcloned into a plasmid vector favorable for DNA preparation and analysis (e.g., commercially available pUC18) to prepare plasmid DNA, from which the cDNA base sequence of the protein of the present invention can be determined by the Maxam Gilbert method (see A.M. Maxam & W. Gilbert, *Proc. Natl. Acad. Sci.*, 74, 560(1977)) or the Sanger method (see F. Sanger & A.R. Coulson, *J. Mol. Biol.*, 94, 441 (1975); and F. Sanger, S. Nicklen & A.R. Coulson, *Proc. Natl. Acad. Sci.*, 74, 5463(1977)).

To determine the base sequence of genomic DNA from the cDNA of the protein of the present invention, for example, plant tissues such as those of leaves, stems and roots are instantaneously frozen with liquid nitrogen, and thoroughly ground with a mortar and pestle or with a whirling blender. From the ground material obtained, genomic DNA is extracted according to an ordinary method as described in Wataru Watanabe (comp.), Masahiro Sugiura (ed.), *Cloning and Sequence (Manual on Plant Biotechnology Experiments)*, Tokyo, Noson Bunka-sha, 1989. The genomic DNA obtained is digested with appropriate restriction enzymes, and the resulting DNA fragments are fractionated by a known method such as sucrose density gradient centrifugation or sodium chloride density gradient centrifugation. For these fractionated DNA fragments, ordinary southern blot hybridization using the cDNA of the protein of the present invention as a probe (genomic southern method) is performed to determine the gene region coding for the desired information.

Further, this gene region is ligated to an appropriate vector such as a commercially available plasmid, phage or cosmid to produce a genomic DNA library. For this library, an ordinary screening by hybridization using the cDNA of the protein of the present invention as a probe is performed to obtain a genomic DNA clone containing a coding region for the amino acid sequence of the protein of the present invention. The genomic DNA clone obtained is subcloned into an appropriate vector favorable for the analysis of a gene sequence, such as a plasmid, and then subjected to the analysis of a base sequence according to an ordinary method, which makes it possible to determine the genomic DNA base sequence containing a coding region for the amino acid sequence of the protein of the present invention. Further, the transcription-initiating site of the genomic DNA of the protein of the present invention can be determined by the primer

extension method as described in M. Bina-Stem et al., *Proc. Natl. Acad. Sci. USA*, 76, 731(1979) or Sollner-Webb & R.H. Reeder, *Cell*, 18, 485(1979), or the SI mapping method as described in A.J. Berk & P.A. Sharp, *Proc. Natl. Acad. Sci. USA*, 75, 1274 (1978). One TATA sequence necessary for the initiation of transcription is present upstream the transcription-initiating site thus determined. In usual cases, a promoter sequence bearing the control of gene expression is present about 1 kb to about 10 kb upstream this transcription-initiating site. The promoter site of the gene of the present invention can be determined by utilizing the fact that the promoter of the gene of the present invention specifically acts in the roots. For example, gene fragments having promoter regions of different lengths are independently ligated to a reporter gene such as GUS, which is then introduced into a certain plant to produce a transgenic plant, and various tissues of the transgenic plant produced are examined for the expression of the reporter gene, by which the promoter site of the gene of the present invention can be finally determined.

On the other hand, a terminator sequence is present in the genomic DNA region corresponding to the site downstream the poly-A sequence which is usually present downstream the poly-A addition signal (i.e., AATAAA and consensus sequence) which is present in the 3' non-translational region downstream the termination codon, and it has an effective function for terminating transcription.

The present invention will be further illustrated by the following examples, which are not to be construed to limit the scope thereof.

#### Example 1: Separation and detection of the protein of the present invention

To 5 g of the root tissue of a commercially available carrot (variety name: Kuroda Gosun) was added 12.5 ml of a buffer for soluble-protein extraction (0.1 M potassium phosphate (pH 8.0), 1 mM EDTA, 0.1% ascorbic acid, 0.25% 2-mercapto-ethanol), and the mixture was ground in an ice bath with a whirling blender (Nihon Seiki) at 2000 rpm for 5 minutes. The ground material was filtered using 4 sheets of gauze, and the filtrate was centrifuged at 10,000 rpm at 4°C for 10 minutes to obtain a supernatant. The supernatant was concentrated to a concentration of about 1 to 2 mg protein/ml by a filter-filtration apparatus (Centriprep-10 (Amicon) and ultra-free CL unit (Millipore)). The concentrate was used as a sample and SDS-polyacrylamide gel electrophoresis was performed on an SDS-polyacrylamide gel with a 10-20% concentration ingredient of acrylamide, "SDS-PAG plate 10/20" (Daiichi Kagaku), in 0.0625 M Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 5% 2-mercaptoethanol, 0.001% bromophenol blue, 1 liter buffer for electrophoresis (0.02 M Tris, 0.192 M glycine, 0.1% SDS, pH 8.4) at a constant current of 60 mA for 60 minutes to separate various proteins on the basis of their molecular weights. The separated proteins were silver-stained using a commercially available kit for silver staining, 2D-silver staining reagent II "Daiichi" (Daiichi Kagaku), and the pattern of protein separation was observed. The results are shown in Figure 1. The protein of the present invention was detected at a position corresponding to a molecular weight of approximately 16 kD, and the ratio of its presence was about 10%, based on the total amount of proteins extracted from the root tissue, as determined by visual observation.

#### Example 2: Determination of the N-terminal amino acid sequence of the protein of the present invention

The concentrate of soluble proteins from the carrot root obtained in Example 1 was separated by the same electrophoresis as described in Example 1, and the separated proteins were stained according to a method as described in P. Matsudaria, *The Journal of Biological Chemistry* (1987). That is, the proteins in the gel were blotted on a commercially available polyvinylidene difluoride (PVDF) membrane in a blotting buffer (10 mM CAPS (3-cyclohexylaminopropane sulfonic acid)-10% methanol (pH 11.0)) at 100 V for 1 hour, stained by 0.1% Coomassie Brilliant Blue (CBB) R250/50% methanol, decolorized in 50% methanol-10% acetic acid, and separated. The band including the protein of the present invention on the PVDF membrane stained by the CBB reagent was cut out by means of a cutter, and the protein of the present invention was collected. Then, 300 pmole of the collected protein of the present invention was analyzed for the amino acid sequence using the amino acid sequencer 473A (Applied Biosynthesis) according to the Edman degradation.

The results are shown in Figure 2. The sequence consisting of 35 amino acids present at the N-terminus of the protein of the present invention was determined.

#### Example 3: Preparation of carrot root cDNA library

In 10 ml of liquid nitrogen was frozen 10 g by fresh weight of the root tissue (9 weeks after seeding) of a commercially available carrot (variety name: Kuroda Gosun), and the frozen tissue was roughly ground by

means of a mallet, and then thoroughly ground with a whirling blender (Nihon Seiki) at 1500 rpm for 25 minutes, while liquid nitrogen was added in several milliliters portions, to give fine powder. After vaporization of the liquid nitrogen, RNA extraction was performed using a commercially available RNA extraction kit (Extract-A-PLANT™ RNA ISOLATION KIT (Clontech)), and the entire RNA was collected from the extract by ethanol precipitation to give 520 µg of the entire RNA. The entire RNA was fractionated in a commercially available oligo-dT column (5'→3') to give 4.8 µg of poly-A RNA. From 1 µg of the poly-A tailed RNA, cDNA synthesis was performed using a commercially available cDNA synthesis kit (cDNA Synthesis System Plus, Amersham) to give 63 ng of cDNA. The cDNA was ligated to an EcoR I-Not I-BamH I adaptor (Takara Shuzo) using commercially available T4 ligase (Takara Shuzo), and a cDNA library composed of λ phage in the XL1-BLUE strain derived from the *Escherichia coli* K-12 strain using EcoR I digested λ ZAPII (Stratagene) and a commercially available *in vitro* packaging kit (GIGA PACK II Gold (Stratagene)).

Example 4: Screening of cDNA clones of the protein of the present invention

Based on the N-terminal amino acid sequence of the protein of the present invention as determined in Example 2, there were synthesized two kinds of synthetic probe DNAs each having the following presumed DNA base sequence:

(1) Probe 1:

5'-GGT GCC CAG AGC CAT GT I CTC GAG ATC ACT TCT TCA  
GTC TCC GCA GAG AAA ATA TTC AGC GGC ATT GTC CTT GAT GTT  
GAT ACA GTT ATT CCC AAG GCT GCC CCC-3' (SEQ ID NO: 3)

(2) Probe 2 (mixed probe):

5'-GGGGCG CAG AGG CAC GTG CTC GAG ATC-3' (SEQ ID NO: 4) and  
5'-GGGGCG CAA AGG CAC GTG CTC GAG ATC-3' (SEQ ID NO: 5)

wherein the symbol "I" in the sequence of probe 1 refers to inosine and probe 2 is a mixture of the above-identified synthetic DNAs. These synthetic probe DNAs were radiolabelled using a commercially available radiolabelling kit (MAGALABEL, Takara Shuzo) to give radiolabelled synthetic probe DNAs.

To *Escherichia coli* XLI-BLUE as a host was added  $5.0 \times 10^4$  pfu of phage cDNA library, and the mixture was incubated at 37°C for 15 minutes and then plated on an NZY plate of ϕ 150 mm (1% NZ amine, 0.5% yeast extract, 0.5% NaCl, 1.5% agar), which was then incubated at 37°C for 7 hours. The plate was cooled to 4°C for 2 hours, and the plaque was transferred on commercially available nylon filters, Hybond-N (Amersham). After air-drying, these filters were denatured successively with an alkali denaturation solution (0.5 M NaOH, 1.5 M NaCl) for 2 minutes, a neutralization solution (0.5 M Tris-HCl, pH 8.0, 1.5 M NaCl) for 2 minutes, and 3 x SSC (0.45 M NaCl, 0.105 M sodium citrate) for 2 minutes, and air-dried, followed by UV-light irradiation for DNA fixation on the filters. In a polyethylene bag were placed ten pieces of the filters, to which 25 ml of prehybridization buffer (6 x SSC (0.9 M NaCl, 0.21 M sodium citrate), 5 x Denhardt's solution (0.1% Ficoll 400, 0.1% polyvinylpyrrolidone, 0.1% BSA), 0.1% SDS, 100 µg/ml denatured salmon sperm DNA) was added, and the contents of the bag were incubated at 45°C for 2 hours. Then, the prehybridization buffer was removed, and 6 ml of fresh prehybridization buffer was added. Further,  $10^7$  cpm of the RI-labelled synthetic probe DNA obtained by the above method was added, and the mixture was incubated at 45°C overnight. The filters were removed and then washed by successive incubation in 50 ml of 6 x SSC at room temperature for 5 minutes; in 100 ml of 6 x SSC at room temperature for 5 minutes; and in 100 ml of 6 x SSC-10% SDS at 45°C for 5 minutes, after which a plaque containing the cDNA for the protein of the present invention was obtained by the RI activity on the filters. The signal plaque obtained was isolated in a solution containing 500 µl of SM (50 mM Tris-HCl, pH 7.5, 0.1 M NaCl, 7 mM MgSO<sub>4</sub>, 0.01% gelatin) and 20 µl of chloroform. Further, to obtain a single clone, *Escherichia coli* XLI-BLUE and  $10^3$ - $10^2$  pfu of phage solution were mixed, and the mixture was incubated at 37°C for 15 minutes and plated on an NZY plate of ϕ 90 mm, which was then incubated at 37°C

overnight. This plate was transferred to nylon membranes by the above method, and two cDNA clones for the protein of the present invention were isolated by plaque hybridization using the same synthetic probe DNA as described above.

- 5 Example 5: Analysis of insert in cDNA clones for the protein of the present invention; and determination of the base sequence coding for the protein of the present invention and the corresponding amino acid sequence

The two cDNA clones obtained in Example 4 were subcloned into plasmid vector pBluescript SK(-) by  
 10 an ordinary method as described in J. Sambrook, E.F. Fritsch & T. Maniatis, *Molecular Cloning Second Edition*, Cold Spring Harbor Laboratory Press (1989) to give plasmid cDNA clones, pC16-1 and pC16-2. The base sequence of the inserts in these cDNA clones was determined by DNA Sequencer 373A (Applied Biosystems) using Taq Dye Primer Cycle Sequencing Kit and Taq Dye Deoxy Terminator Cycle Sequencing Kit (Applied Biosystems). The pC16-1 and pC16-2 both have an insert having the same base  
 15 sequence. For this reason, these plasmids were identical and designated pC16 (see Figure 3). The pC16 has been deposited as *Escherichia coli* XL1-BLUE/pC16 (accession number: FERM BP-4469) with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology under the Budapest treaty (date of the original deposit: November 17, 1993). The insert had a size of 739 bases (see Sequence Listing, SEQ ID NO: 6), in which 465 bases at the 14th to the 478th positions were  
 20 found to constitute a coding region for the 16 kD protein which can be specifically expressed in carrot roots. It was also found that bases at the 479th to the 717th positions constitute a 3' non-translational region and twenty-two pieces of adenine at the 718th to the 739th positions constitute a poly-A tail. From the base sequence of the coding region for the 16 kD protein which can be specifically expressed in carrot roots, the entire amino acid sequence of the protein was determined (see Sequence Listing, SEQ ID NO: 2). The  
 25 protein of the present invention comprises 154 amino acids (see Sequence Listing, SEQ ID NO: 1), and it has a calculated molecular weight of 16125.02.

Example 6: Tissue-specific expression of the protein of the present invention

The seeds of a commercially available carrot (variety name: Kuroda Gosun) were sowed in the soil of a pot in a greenhouse, and cultivated. On the 4th, 7th, 9th and 21st weeks after the sowing, the plants were  
 30 collected and separated into the root (subterranean) tissue and the leaf (terrestrial) tissue. All the proteins in the respective tissues were extracted and concentrated by the same method as described in Example 1, and SDS-polyacrylamide gel electrophoresis was performed with 1 µg of the protein per lane as a sample under the same conditions as described in Example 1, by which various proteins were separated on the basis of their molecular weights and compared for the expression of the protein of the present invention.

35 The results are shown in Figure 4. All the proteins in the entire plant on the 4th week was compared with those in the root and leaf at the 7th, 9th and 21st weeks, respectively, and it is found that the expression of the protein of the present invention is notably different. That is, the protein of the present invention was not expressed in the young plants or in the respective leaf tissues, but expressed only in the root tissues; it is further clear that the protein of the present invention is a reserve protein having an  
 40 increased ratio to all the proteins with a progress of the growth of the root tissues.

Example 7: Comparison between the varieties of the protein of the present invention

From the root tissues of various commercially available carrots (variety names: Kuroda Gosun, Early  
 45 Chantenee, Imperator and Nuntesscarlet), all the proteins were extracted and concentrated by the same method as described in Example 1, and SDS-polyacrylamide electrophoresis was performed with 1 µg of the protein per lane as a sample under the same conditions as described in Example 1, by which various proteins were separated on the basis of their molecular weights and compared for the expression of the protein of the present invention.

50 The results are shown in Figure 5. It was found that the protein of the present invention is the most important protein corresponding to about 10% of all the proteins, regardless of the kind of variety.

Example 8: Homology of the protein of the present invention

55 Genes and amino acid sequences, which have high homology with the base sequence coding for the protein of the present invention and the corresponding amino acid sequence, respectively, as determined in Example 5, were searched on the data bases EMBL and NBRF. As a result, it was found that the gene and amino acid sequence of the present invention exhibit the highest homology with the genes and amino acid

sequences of several proteins related to plant disease resistance (see Figure 6). The proteins having the highest homology were Pathogenesis-related Protein (PR-Protein) 1-3, 1-1 and 1-2 of parsley belonging to the family *Umbelliferae* similarly to carrot (see Figure 6). Relatively high homology was also observed in the mRNA of 3T3-7, 3T3-9 and 3T3-M1 clones of HSP60 which is a heat shock protein of mice.

As described above, the gene and plasmid of the present invention are useful for higher expression of the protein of the present invention, which is specifically present in carrot roots, and it becomes possible to develop high protein carrots by such higher expression.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: Sumitomo Chemical Co., Ltd.  
 (B) STREET: 5-33, Kitahama 4-chome, Chuo-ku  
 (C) CITY: Osaka-shi, Osaka-fu  
 (E) COUNTRY: Japan  
 (F) POSTAL CODE (ZIP): none

(ii) TITLE OF INVENTION: CARROT 16 KD PROTEIN, GENE CODING FOR SAID PROTEIN AND PLASMID CONTAINING SAID GENE

(iii) NUMBER OF SEQUENCES: 15

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
 (B) COMPUTER: IBM PC compatible  
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS  
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: JP 327943/1993  
 (B) FILING DATE: 24-DEC-1993

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 154 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: *Daucus carota* L. (carrot; variety name: Kuroda Gosun)  
 (F) TISSUE TYPE: Root (subterranean part)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met	Gly	Ala	Gln	Ser	His	Ser	Leu	Glu	Ile	Thr	Ser	Ser	Val	Ser	Ala
1				5					10					15	
Glu	Lys	Ile	Phe	Ser	Gly	Ile	Val	Leu	Asp	Val	Asp	Thr	Val	Ile	Pro
			20					25					30		
Lys	Ala	Ala	Pro	Gly	Ala	Tyr	Lys	Ser	Val	Asp	Val	Lys	Gly	Asp	Gly
			35				40					45			
Gly	Ala	Gly	Thr	Val	Arg	Ile	Ile	Thr	Leu	Pro	Glu	Gly	Ser	Pro	Ile
			50			55					60				



Thr Ser Met Thr Val Arg Thr Asp Ala Val Asn Lys Glu Ala Leu Thr  
 65 70 75 80  
 Tyr Asp Ser Thr Val Ile Asp Gly Asp Ile Leu Leu Glu Phe Ile Glu  
 85 90 95  
 Ser Ile Glu Thr His Met Val Val Val Pro Thr Ala Asp Gly Gly Ser  
 100 105 110  
 Ile Thr Lys Thr Thr Ala Ile Phe His Thr Lys Gly Asp Ala Val Val  
 115 120 125  
 Pro Glu Glu Asn Ile Lys Phe Ala Asp Ala Gln Asn Thr Ala Leu Phe  
 130 135 140  
 Lys Ala Ile Glu Ala Tyr Leu Ile Ala Asn  
 145 150

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 465 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA to mRNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Daucus carota L. (carrot; variety name:  
 Kuroda Gosun)  
 (F) TISSUE TYPE: root (subterranean part)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

ATGGGTGCCC AGAGCCATTC ACTCGAGATC ACTTCTTCAG TCTCCGCAGA GAAAATATTC 60  
 AGCGGCATTG TCCTTGATGT TGATACAGTT ATTCCCAAGG CTGCCCCCGG AGCTTACAAG 120  
 AGTGTGCATG TTAAAGGAGA CGGTGGAGCT GGAACCGTCA GAATTATCAC CCTTCCCGAA 180  
 GGTAGCCCAA TCACCTCAAT GACGGTTAGG ACTGATGCAG TGAACAAGGA GGCCTTGACA 240  
 TACGATTCCA CAGTCATTGA TGGAGACATC CTTCTAGAAT TCATCGAATC CATTGAAACC 300  
 CATATGGTAG TTGTGCCAAC TGCTGACGGA GGTAGCATT CCAAGACCAC TGCCATATTC 360  
 CACACCAAAG GCGATGCCGT GGTTCCTGAG GAGAACATCA AGTTTGCAGA TGCTCAGAAC 420  
 ACTGCTCTTT TCAAGGCTAT TGAGGCCTAC CTCATTGCTA ATTAA 465

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION: 18

(D) OTHER INFORMATION: /note= "n at position 18 refers to inosine."

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGTGCCCGAGA GCCATGTNCT CGAGATCACT TCTTCAGTCT CCGCAGAGAA AATATTCAGC 60

GGCATTGTCC TTGATGTTGA TACAGTTATT CCCAAGGCTG CCCCC 105

15

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

20

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGGGCGCAGA GGCACGTGCT CGAGATC 27

(2) INFORMATION FOR SEQ ID NO: 5:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

40

GGGGCGCAAA GGCACGTGCT CGAGATC 27

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 739 base pairs

45

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

50

55

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: *Daucus carota* L. (carrot; variety name:  
Kuroda Gosun)

(F) TISSUE TYPE: root (subterranean part)

5

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

10 CATTCTAAAT ATCATGGGTG CCCAGAGCCA TTCACTCGAG ATCACTTCTT CAGTCTCCGC 60  
 AGAGAAAATA TTCAGCGGCA TTGTCCTTGA TGTGATACA GTTATTCCCA AGGCTGCCCC 120  
 CGGAGCTTAC AAGAGTGTG ATGTTAAAGG AGACGGTGGA GCTGGAACCG TCAGAATTAT 180  
 15 CACCCTTCCC GAAGGTAGCC CAATCACCTC AATGACGGTT AGGACTGATG CAGTGAACAA 240  
 GGAGGCCTTG ACATACGATT CCACAGTCAT TGATGGAGAC ATCCTTCTAG AATTCATCGA 300  
 ATCCATTGAA ACCCATATGG TAGTTGTGCC AACTGCTGAC GGAGGTAGCA TTACCAAGAC 360  
 CACTGCCATA TTCCACACCA AAGGCGATGC CGTGGTTCCT GAGGAGAACA TCAAGTTTGC 420  
 20 AGATGCTCAG AACACTGCTC TTTTCAAGGC TATTGAGGCC TACCTCATTG CTAATTAAGC 480  
 TGAGCTCTCA ACTTCCGTAA TTTTATGAGT GAGTGGAGGA ATTGCAACGT TTTCTTTTGT 540  
 GTTTTGT TTTT CGAGCAACTT CATAATTTAC AGAGTGAGTG ACAGTCAGTG ACAGAATTGC 600  
 25 AACTTTCTCT TTGTACTTTG TTGTGACTTG TGATGAATAA CTTCATCTGG CTGGTAATGT 660  
 ATGCGATCTT TTTAAATAAT ATGCACTATT ATTAAACCAA TAATCATATT CATTCTCAAA 720  
 AAAAAAAAAA AAAAAAAAAA 739

30

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

## (ii) MOLECULE TYPE: peptide

40

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Gly Ala Gln Ser His Ser Leu Glu Ile Thr Ser Ser Val Ser Ala Glu  
 1 5 10 15  
 Lys Ile Phe Ser Gly Ile Val Leu Asp Val Asp Thr Val Ile Pro Lys  
 20 25 30  
 Ala Ala Pro  
 35

50

55

## (2) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Gly Ala Gln Ser His Ser Leu Glu Ile Thr Ser Ser Val Ser Ala  
 1 5 10 15  
 Glu Lys Ile Phe Ser Gly Ile Val Leu Asp Val Asp Thr Val Ile Pro  
 20 25 30  
 Lys Ala Ala Pro Gly Ala Tyr Lys Ser Val Asp  
 35 40

## (2) INFORMATION FOR SEQ ID NO: 9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Val Lys Gly Asp Gly Gly Ala Gly Thr Val Arg Ile Ile Thr Leu Pro  
 1 5 10 15

## (2) INFORMATION FOR SEQ ID NO: 10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Gly Val Gln Lys Ser Glu Val Glu Ala Thr Ser Ser Val Ser Ala  
 1 5 10 15

Glu Lys Leu Phe Lys Gly Leu Cys Leu Asp Ile Asp Thr Leu Leu Pro  
                   20                                  25                                  30  
 Arg Val Leu Pro Gly Ala Ile Lys Ser Ser Glu Thr Leu Glu Gly Asp  
                   35                                  40                                  45  
 Gly Gly Val Gly Thr Val Lys Leu Val His Leu Gly  
                   50                                  55                                  60

## (2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 60 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Met Gly Val Gln Lys Ser Glu Val Glu Thr Thr Ser Ser Val Ser Ala  
   1                                  5                                  10                                  15  
 Glu Lys Leu Phe Lys Gly Leu Cys Leu Asp Ile Asp Thr Leu Leu Pro  
                   20                                  25                                  30  
 Gln Val Leu Pro Gly Ala Ile Lys Ser Ser Glu Thr Leu Glu Gly Asp  
                   35                                  40                                  45  
 Gly Gly Val Gly Thr Val Lys Leu Val His Leu Gly  
                   50                                  55                                  60

## (2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 60 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Glu Gly Ser Pro Ile Thr Ser Met Thr Val Arg Thr Asp Ala Val Asn  
   1                                  5                                  10                                  15  
 Lys Glu Ala Leu Thr Tyr Asp Ser Thr Val Ile Asp Gly Asp Ile Leu  
                   20                                  25                                  30  
 Leu Glu Phe Ile Glu Ser Ile Glu Thr His Met Val Val Val Pro Thr  
                   35                                  40                                  45

Ala Asp Gly Gly Ser Ile Thr Lys Thr Thr Ala Ile  
50 55 60

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 60 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Asp Ala Ser Pro Phe Lys Thr Met Lys Gln Lys Val Asp Ala Ile Asp  
1 5 10 15  
Lys Ala Thr Phe Thr Tyr Ser Tyr Ser Ile Ile Asp Gly Asp Ile Leu  
20 25 30  
Leu Gly Phe Ile Glu Ser Ile Asn Asn His Phe Thr Ala Val Pro Asn  
35 40 45  
Ala Asp Gly Gly Cys Thr Val Lys Ser Thr Ile Ile  
50 55 60

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 35 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Phe His Thr Lys Gly Asp Ala Val Val Pro Glu Glu Asn Ile Lys Phe  
1 5 10 15  
Ala Asp Ala Gln Asn Thr Ala Leu Phe Lys Ala Ile Glu Ala Tyr Leu  
20 25 30  
Ile Ala Asn  
35

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 35 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Phe Asn Thr Lys Gly Asp Ala Val Val Pro Glu Glu Asn Ile Lys Phe  
 1 5 10 15  
 10 Ala Asn Asp Gln Asn Leu Thr Ile Phe Lys Ala Val Glu Ala Tyr Leu  
 20 25 30  
 Ile Ala Asn  
 35

15

### Claims

- 20 1. A gene encoding the protein having the following features:  
 (a) it is specifically expressed in carrot roots; (b) it has a molecular weight of 16kD; and (c) it corresponds to about 10% of root tissue protein.
- 25 2. The gene according to claim 1, wherein said protein comprises 154 amino acids.
3. The gene according to claim 1 or 2, wherein said protein has the amino acid sequence shown in Seq. ID No. 1.
- 30 4. The gene according to any one of claims 1 to 3, which has the base sequence as shown in Seq. ID No. 2.
5. A vector comprising the gene according to any one of claims 1 to 4.
- 35 6. The vector according to claim 5 which is an expression vector.
7. The vector according to claim 5 or 6, which is a plasmid.
8. A host transformed with the vector according to any one of claims 5 to 7.
- 40 9. The host according to claim 8 which is a carrot or another root vegetable crop such as radishes, turnips, sugar beets and burdocks, or Escherichia coli, preferably E. coli XL1-BLUE/pC16 (FERM BP-4469).
- 45 10. A protein encoded by the gene of any one of claims 1 to 4 or expressed by the vector of claim 5 or 6 or by the host of claim 8 or 9.
11. A process for producing the protein of claim 10, which comprises culturing the host of claim 8 or 9 under suitable conditions and recovering the produced protein.

50

55




Fig. 1



N-terminus

1 35  
GAQSHSLEITSSVSAEKIFSGIVLDVDTVIPKAAP

(SEQ ID NO: 7)

Fig. 2




Fig. 3




Fig. 4





Fig. 5

BNSDOCID: <EP\_\_\_0659884A2\_I\_>



1

2



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 659 884 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
06.08.1997 Bulletin 1997/32

(51) Int. Cl.<sup>6</sup>: **C12N 15/29**, C12N 1/21,  
C07K 14/415

(43) Date of publication A2:  
28.06.1995 Bulletin 1995/26

(21) Application number: **94120679.9**

(22) Date of filing: **27.12.1994**

(84) Designated Contracting States:  
**DE FR GB**

(30) Priority: **24.12.1993 JP 327943/93**

(83) Declaration under Rule 28(4) EPC (expert  
solution)

(71) Applicant: **SUMITOMO CHEMICAL COMPANY  
LIMITED**  
**Osaka-shi, Osaka 541 (JP)**

(72) Inventors:

- **Yamamoto, Mika**  
**Sakado-shi, Saitama-ken (JP)**
- **Oheda, Kenji,**  
**Pereniarufulakksu 1,6**  
**Kyoto-shi, Kyoto-fu (JP)**

(74) Representative: **VOSSIUS & PARTNER**  
**Postfach 86 07 67**  
**81634 München (DE)**

(54) **Carrot 16 kd protein, gene coding for said protein and plasmid containing said gene**

(57) There are disclosed a protein corresponding to a molecular weight of approximately 16 kD, which can be specifically expressed in carrot roots, a gene coding for the protein, and a plasmid containing the gene.

**EP 0 659 884 A3**



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 94 12 0679

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	PROTOPLASMA, (1989) VOL. 150, NO. 2-3, PP. 139-149. , XP000672545 SLAY, R. M. ET AL: "Plasma membrane proteins associated with undifferentiated and embryonic Daucus carota tissue" * the whole document *	1,2	C12N15/29 C12N1/21 C07K14/415
A	PLANT PHYSIOLOGY, vol. 100, 1992, pages 374-381, XP002031535 TURANO, F.J., ET AL.: "Identification and expression of a cDNA clone encoding aspartate aminotransferase in carrot" * the whole document *	1,2	
A	PLANT PHYSIOLOGY, (1988) VOL. 88, NO. 2, PP. 444-449. , XP000672546 GRIMES, H. D. ET AL: "Plant plasma membrane proteins. II. Biotinylation of Daucus carota protoplasts and detection of plasma membrane polypeptides after SDS-PAGE"	1,2	
A	THE PLANT CELL, vol. 2 , 1990, pages 457-467, XP002031536 HATZOPOULOS, P., ET AL.: "Interaction of nuclear factors with upstream sequences of a lipid body membrane protein gene from carrot" * the whole document *	1,2	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C12N C07K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 4 June 1997	Examiner Maddox, A
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 150 03.82 (P0401)





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 94 12 0679

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	PLANT SCI (LIMERICK) 88 (1). 1993. 1-11. , XP000675018 ROSS J H E ET AL: "DIFFERENTIAL ACCUMULATION OF STORAGE PRODUCTS IN DEVELOPING SEEDS AND SOMATIC CELL CULTURES OF DAUCUS-CAROTA L." * the whole document *	1,2	
A	NUCLEIC ACIDS RESEARCH, vol. 18, 1990, page 2826 XP002031552 ULRICH, T.U., ET AL.: "Sequence of EMB-1, an mRNA accumulating specifically in embryos of carrot" * the whole document *	1,2,5,8	
A	MOL. GEN. GENET., vol. 213, 1988, pages 93-98, XP000673889 SOMSSICH, I.E., ET AL.: "Gene structure and in situ transcript localization of pathogenesis-related protein 1 in parsley" * the whole document *	1,2	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 4 June 1997	Examiner Maddox, A
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 01.82 (P04C01)

**THIS PAGE BLANK (USPTO)**